



Research



FERAL CATS AND THE PUBLIC: A HEALTHY RELATIONSHIP

Public health policies all over the country reflect the scientific evidence: feral cats live healthy lives outdoors and don't spread disease to people. But advocates of catch and kill programs continue to justify this cruel practice by insisting that feral cats represent a threat to public health and that they do spread disease. "There's simply no evidence to back up these claims," says Deborah L. Ackerman, M.S., Ph.D., an adjunct associate professor of epidemiology at the UCLA School of Public Health.

More and more, public health officials are embracing Trap-Neuter-Return for feral cats and replacing outdated policies based on unfounded fears.

"I'm not a cat or animal lover," says Ron Cash, director of the Atlantic City Department of Health and Human Services, "But I believe the complaints and hysteria about disease as result of feral cats are overblown."



Most diseases that infect cats can only be spread from cat to cat, not from cat to human. You are much more likely to catch an infectious disease from the person standing in line with you at the grocery store than from a cat.¹ In fact, a 2002 review of cat-associated diseases published in the Archives of Internal Medicine concluded that, "cats should not be thought of as vectors for disease transmission."²

Infectious diseases can only spread from cats to humans via direct contact with either the cat or its feces, and feral cats typically avoid humans. Statistics from the Centers for Disease Control and Prevention (CDC) show that cats are rarely a source of disease, and that it is unlikely for anyone to get sick from touching or owning a cat.³ "Feral cats pose even less risk to public health than pet cats because they have minimal human contact, and any contact that does occur is almost always initiated by the person," says Ackerman.

Science shows feral cat colonies aren't a risk to humans

The health risks that catch and kill advocates most often blame on cats are intestinal parasites, rabies, flea-borne typhus, and toxoplasmosis. Yet the spread of these diseases has never been conclusively linked to feral cats.

Parasites are Species Specific

Ackerman says that the risk of catching an intestinal parasite like *Cryptosporidium* and *Giardia* from cats has been vastly over-hyped. Molecular studies show that these parasites are usually species specific—meaning that the type that infects cats does not infect humans—and “some studies even suggest that cats and other animals are more likely to catch these parasites from humans than vice-versa,” according to Ackerman. Feral cats live full, healthy lives outdoors—there is no reason for them to be killed in shelters.

No Danger from Rabies

The notion that stray cats spread rabies is another empty argument used by advocates of catch and kill programs, says Ackerman. The last confirmed cat-to-human transmission of rabies occurred in 1975. The risk of catching rabies from a feral cat is almost non-existent. Statistics from the CDC show that as a source of rabies infections, cats rank way behind wild animals like bats, skunks, and foxes who account for more than 90 percent of reported cases of the disease.⁴

Trap-Neuter-Return is a safeguard against rabies, because “the vaccination component of TNR programs ensures that the cats in managed colonies cannot catch or spread rabies,” says Ackerman.

Even in the unlikely event that a feral cat develops rabies, it can't spread the disease to people without biting them, and feral cats rarely seek direct contact with humans. The idea that cats will unexpectedly jump out of alleys and bite children is just as ridiculous as it sounds. A 1998 analysis showed that about 90 percent of cat bites were provoked, and the vast majority of cat bites are caused by pets.⁵

Since Atlantic City, New Jersey began its TNR program in 2001, the city hasn't had a single complaint about feral cat bites or scratches. Learn more about why feral cats do not spread rabies at www.alleycat.org/PublicHealthVictory.

Flea-borne Typhus is Rare and Cats Don't Play a Part in Flea Arrival or Growth

Flea-borne typhus is another infectious disease sometimes erroneously blamed on feral cats. The disease is caused by *Rickettsia* bacteria that infect fleas, and most U.S. cases occur in Texas, Hawaii, and California. Although infected fleas may hitch a ride on feral cats, the chance of becoming infected with flea-borne typhus via a feral cat is extremely low. Ackerman says, “flea-borne typhus is rare even in areas such as Southern California, where the disease is endemic.” For instance, in 2009, Orange County, California reported 12 cases of flea-borne typhus out of a population of 3 million residents⁶, making the chance of infection just 1 in 250,000—about the same as the risk of being hit by an asteroid.⁷

Removing cats does not halt the spread of flea-borne typhus, because cats don't spread the disease—the fleas themselves do. Cats are merely a host for fleas and if the cats are eliminated, the fleas simply find another host like squirrels and raccoons. “Fleas are very versatile. They live on cats, dogs, opossums, rats, and mice,” Ackerman says.

For this reason, public health officials in Texas, where flea-borne typhus is endemic, have focused their efforts on controlling fleas rather than their hosts. Outbreaks are rarely traced to cats. In 2008, the CDC and Texas health authorities examining a cluster of flea-borne typhus in Austin found the *Rickettsia* bacteria in only 18 percent of cats, as compared to 44 percent of dogs and 71 percent of opossums, near the homes of people infected with the disease.⁸

Most Cases of Toxoplasmosis Stem from Undercooked Food, Not Cats

Catch and kill advocates argue that feral cats transmit toxoplasmosis, a parasitic disease that spreads via *Toxoplasma* oocysts shed in the feces of an infected animal. But studies show that the overwhelming majority of toxoplasmosis cases actually result from eating undercooked meat. According to CDC statistics, toxoplasmosis is the third leading cause of food-borne illness-related death in the U.S.⁹

Pregnant women and their fetuses face a higher risk from the disease—a fact that catch and kill advocates often abuse to incite public paranoia. But a study published in the *Archives of Internal Medicine* in 2002 concluded that pregnant women were unlikely to catch toxoplasmosis from a cat.¹⁰

It's rare for anyone to catch toxoplasmosis from a household pet (cats are not the only carriers; dogs, birds, and other mammals can also carry the parasite), let alone a feral cat with whom they have no contact. Even if a cat is infected with *Toxoplasma*, it typically only sheds the disease-spreading oocysts for a few weeks. To catch an infection, a person would need to have direct contact with these infected feces. Most people go out of their way to avoid touching the contents of their pet cat's litter box, and they're even less likely to touch feral cat feces. In other words, even if a feral cat leaves feces in your garden, you would need to touch it and then somehow ingest the feces to get toxoplasmosis.

Colony Caregivers are as Healthy as Everyone Else

Maybe the best proof that feral cats are not a health risk is that feral cat caregivers are healthy. “If feral cats transmitted disease to humans,” says Ackerman, “colony caregivers, who spend more time around feral cats than most people, would experience a heightened rate of disease, and this simply isn't the case.” None of the many caregivers she's interviewed have ever reported becoming sick from their work with feral cats. No study has ever shown that colony caregivers have any increased risk of disease, despite their regular contact with feral colonies.

Catch and Kill Doesn't Improve Public Health

“Catch and kill policies are fear-based and rely on old wives' tales and flawed research to justify prejudice against cats,” says Ackerman. Removing feral cats is not a sustainable solution, because that only opens up new territory for other feral cats to use.

Learn more about this vacuum effect at www.alleycat.org/VacuumEffect. According to Ackerman, there's absolutely no evidence that catch and kill policies reduce the incidence of human disease.

Trap-Neuter-Return Programs Protect Public Health and Prevent the Spread of Disease

Trap-Neuter-Return programs stabilize feral cat populations, and the vaccination component ensures that cats are protected against disease. These programs also allow cat caregivers and public health officials to monitor the health of cats and ensure that they're immunized—and that “protects the health of cats and humans alike,” says Cash. Catch and kill programs offer no such benefits, because cats are simply removed without regard to their health.

“TNR is good public health policy,” says Cash. Atlantic City has been collaborating with Alley Cat Allies since 2001 to manage feral cat colonies under the city’s famous boardwalk. The TNR program that Atlantic City developed with Alley Cat Allies has never posed any health problems to the community, says Cash.

“Before our relationship with Alley Cat Allies, I was getting numerous complaints about feral cats,” he said. But since Alley Cat Allies began managing these colonies with TNR, the problems have ceased entirely. “The [cat] population that’s here is much healthier,” says Cash. “They’re coexisting with people very well now. Most people don’t even know the cats are there.”

While catch and kill advocates cling to outdated thinking and hyped-up stories, the people studying, teaching, and defending public health recognize that feral cats do not spread disease to people. Policies based on fear, hype, and hysteria serve neither the public nor the cats and will only end in more cats being killed.

Instead, feral cat policies should reflect the science and the facts—feral cats are healthy animals. From a public health standpoint as well as a humane one, the best approach for feral cats is Trap-Neuter-Return because it benefits the cats and the community.

¹ American Association of Feline Practitioners and the Cornell Feline Health Center, Cornell University, College of Veterinary Medicine. *Zoonotic Disease: What Can I Catch From My Cat?* 2002.

² Kravetz, Jeffrey D., and Daniel G. Federman. "Cat-Associated Zoonoses." *Arch. Intern Med* 162, no. 17 (2002): 1945-1952.

³ Centers for Disease Control and Prevention. *Diseases from Cats*. July 28, 2010. <http://www.cdc.gov/healthypets/animals/cats.htm> (accessed October 25, 2010).

⁴ Centers for Disease Control and Prevention. *Rabies - Epidemiology*. September 18, 2007. <http://www.cdc.gov/rabies/epidemiology.html>.

⁵ Patrick, G.R., and KM O'Rourke. "Dog and Cat Bites: Epidemiologic Analyses Suggest Different Prevention Strategies." *Public Health Report*, 1998: 252-257.

⁶ *Notifiable Diseases in Animals: Joint Meeting of the CCLHO Communicable Disease Control and Environmental Health Committees*, April 15 (2010) (written statement of Deborah L. Ackerman, M.S., Ph.D., Adjunct Associate Professor of Epidemiology, UCLA School of Public Health on Free-Roaming Cats and the Public Health).

⁷ Britt, Robert Roy. *The Odds of Dying*. January 5, 2005. http://www.livescience.com/environment/050106_odds_of_dying.html (accessed October 25, 2010).

⁸ Adjemian, Jennifer, et al. "Murine Typhus in Austin, Texas, USA, 2008." *Emerging Infectious Diseases*, 2010: 412-417.

⁹ Centers for Disease Control and Prevention. *Toxoplasmosis*. January 11, 2008. <http://www.cdc.gov/toxoplasmosis/> (accessed October 25, 2010).

¹⁰ American Association of Feline Practitioners and the Cornell Feline Health Center, Cornell University, College of Veterinary Medicine. *Zoonotic Disease: What Can I Catch from My Cat?* 2002.

ABOUT ALLEY CAT ALLIES

Alley Cat Allies is the only national nonprofit dedicated exclusively to the protection and humane treatment of all cats.

For over 25 years, Alley Cat Allies has led the movement to save cats' lives. With advocacy, education, and grassroots organizing, we champion innovative programs and policies that serve communities and save cats. Since our founding in 1990, we have popularized Trap-Neuter-Return (TNR) as the only humane and effective approach to community cats. Before Alley Cat Allies, TNR was virtually unknown in America, and no local governments supported it. Today, more than 600 municipalities have officially embraced TNR.

Alley Cat Allies has changed how all cats are treated—pet, stray, feral, and each cat in between. We advocate for a variety of progressive policies to protect cats, including the transformation of the American animal shelter system, which still routinely kills 70 percent of all cats who enter.

Thanks to our more than half a million supporters and countless advocates, Alley Cat Allies works in hundreds of communities each year. To learn more about how we save cats' lives, visit www.alleycat.org.